

Immune Deficiency Immunoglobulin Therapy								
То	From			Number of Pages including Cover				
ntake Phone			Phone	Fax				
Patient Name			DOB Date					
Allergies			Height Weight					
Rx: Intravenous Route   IVIGgrams daily forday(s). OR IVIGgrams/kilogram daily given overnon-consecutive day(s).   Repeat course everyweek(s) for a total ofcourse(s). Dose will be rounded to nearest vial size.   Rx: Subcutaneous Route								
IG grams each month given as doses OR IG grams times per month. Administer SQIG using sites at a time. Repeat week(s). Ok to round dose to nearest vial size. Refill x 1yr.								
Diagnosis	ICD-9	ICD-10	Diagnosis			ICD-9	ICD-10	
Common Variable Immunodeficiency with Predominant Immunoregulatory T-Cell Disor	ders 279.10	D83.1	Selective deficier	icy of Immunogl	276.02	D80.4		
Wiskott-Aldrich Syndrome	279.12 Da		Selective deficier G [IgG] Subclasse		279.03	D80.3		
Combined Immunodeficiency, Unspecified		D81.9	Hereditary Hypog		279.04	D80.0		
Severe Combined Immunodeficiency [SCID] with Low T- and B- Cell Numbers	279.2	D81.1	Immunodeficienc	y with Increased	279.05	D80.5		
Severe combined Immunodeficiency [SCID] with Low or Normal B-Cell Numbers		D81.2	Other Common Ve	ariable Immunoo	279.06	D83.8		
Selective deficiency of Immunoglobulin A [I	Selective deficiency of Immunoglobulin A [IgA] 279.01 D80.2			Common Variable Immunodeficiency, Unspecified			D83.9	
Other:								
IV Access Device Peripheral Central Hydration: Infuse 500 ml of NS. Start at 100ml/hr 30 minutes before the IVIG starts; then run concurrently with IVIG infusion.								
Medi-Cal ID# Refill x 1Year If applicable, flush intravenous access device per Hom					device per Home C	are Services p	protocol:	
Per Home Care Services recommendation:			Access		NS	Heparin 100 u/ml		
-ACETAMINOPHEN 650 MG (325mg X 2) orally PRE-IVIG			Peripheral		fore/after use	1-3 ml		
-DIPHENHYDRAMINE 25 MG orally PRE-IVIG None			Midline, Central (Non-Port), Pl		efore/after use 3 fter blood draw after		าไ	
Other premed orders: Other premed orders:			Implanted Port		efore/after use fter blood draw	5 ml after last NS		
Other premed orders: Epi-Pen 0.3mg 2-Pak Auto-Injector			Groshong PICC, Midli		efore/after use fter blood draw			
If needed, a Basic Metabolic Panel (Chem 7) to be drawn prior to first dose. If necessary Patient may have first dose administered in the home by a skilled nurse, unless instructed otherwise. Pharmacy to provide all equipment and medical supplies necessary to administer therapy as appropriate.								
Prescriber Signature:	Date							
Print Prescriber Name			NPI#					
Please fax the following information: Immunoglobulin order – include dose, route of administration, frequency, duration, and any pre-medications OR use prescription order section above Patient demographics – include insurance information. <b>We will obtain authorization</b> unless the insurance dictates otherwise H & P <b>OR</b> progress note(s) describing diagnosis and clinical status Labs – BUN/Creatinine (preferred within last 90 days), Immunoglobulin Panel								
I authorize Vital Care Infusion Services LLC and its representatives to initiate any insurance prior authorization process I order. I understand that I can revoke this designation at any time by providing written notice to Vital Care. Date:								

PRESCRIBER MUST MANUALLY SIGN - STAMP SIGNATURE, SIGNATURE BY OTHER PERSONNEL AND COMPUTER-GENERATED SIGNATURES WILL NOT BE ACCEPTED

The attached document(s) contain confidential information which may be considered to be Protected Health Information and therefore required to be maintained as private and secure under HIPAA. The documents may also contain information which is otherwise considered to be privileged under state and federal laws. This communication is for the intended recipient only. If you are not the intended recipient, or a person responsible for delivering this communication to the intended recipient, you are prohibited from viewing, copying and/or distributing the information contained herein. Unlawful disclosure of the information attached may subject you to monetary penalties and sanctions. If you have received this communication in error, you should notify the sender immediately and thereafter permanently destroy all copies of this document in its entirety.

This form is not considered an order or prescription for medical services and/or supplies unless and until it is formally authorized by a healthcare provider in compliance with applicable laws and regulations.